

26 **Effect of waterlogging on soil biochemical properties and organic matter quality**
27 **in different salt marsh systems**

28

29 Chiara Ferronato^a, Sara Marinari^b, Ornella Francioso^a, Diana Bello^c, Carmen Trasar-Cepeda^c, Livia
30 Vittori Antisari^a

31

32 ^a Dipartimento di Scienze e Tecnologie Agro-Alimentari, Alma Mater Studiorum – Università di
33 Bologna, Via Fanin 44, 40127 Bologna, Italy

34 ^b Dipartimento per l' Innovazione dei Sistemi Biologici, Agroalimentari e Forestali, Università della
35 Tuscia, Via S. Camillo de Lellis, 01100 Viterbo, Italy

36 ^c Departamento de Bioquímica del Suelo, Instituto de Investigaciones Agrobiológicas de Galicia-
37 Consejo Superior de Investigaciones Científicas, Apartado 122, 15780 Santiago de Compostela,
38 Spain

39

40 **Abstract**

41 This study investigated the effects of hydroperiod on soil organic matter quality in three different salt
42 marshes in the Baiona lagoon (N Italy) representing terrestrial, intertidal and subaqueous ecosystems
43 in the area. The study specifically aimed to gain some insight into how soil waterlogging
44 (hydroperiod) affects the chemical and biological properties of soils as well as the quality and
45 structure of the soil organic matter (SOM). Total contents of selected nutrients, total organic carbon
46 and carbon stable isotope (δC^{13}) were measured in all soil profiles. The results of these analyses
47 enabled us to define the different origin of the SOM by discriminating between terrestrial and aquatic
48 SOM sources. The findings also show that accumulation of nutrients and SOM is significantly
49 magnified in intertidal systems, in which pedoturbation effects induced by water movements are
50 particularly strong. In addition, DRIFT spectra of humic acids revealed the changes in the main

functional groups in relation to increased waterlogging, highlighting the lower aromaticity and complexity in subaqueous soils (SASs), which is possibly due to the effect of the soil water saturation on the chemical and biological SOM transformation processes. Microbial biomass carbon (MBC), microbial quotient (Q_{mic}) and the activities of some soil enzymes were measured to estimate soil metabolic activity in the systems and to evaluate how the microbial pool contributes to transforming the SOM. In all systems, the enzymatic activities were generally higher in subsurface horizons than in the surface horizon. This unexpected behaviour can be explained by the combined effect of water movement, erosion processes and preservation of SOM under anaerobic conditions. This study represents an attempt to investigate and understand the ongoing degradation processes in salt marsh ecosystems. The findings emphasize the strong influence of water flow and erosional processes associated with soil waterlogging on chemical and biological reactions in intertidal and subaqueous systems.

Keywords: Subaqueous soils; Soil organic matter transformations; Soil waterlogging; Enzyme activity; Salt marshes

1. Introduction

Salt marshes and wetlands represent some of the largest and most efficient C sink systems in the world (the so called “blue carbon”). It has been estimated that wetland systems hold 45-70% of all terrestrial C (Mitra et al., 2005; Morrissey et al., 2014) and that salt marshes can store about $87.2 \pm 9.6 \text{ Tg C yr}^{-1}$ in comparison with the $53 \pm 9.6 \text{ Tg C yr}^{-1}$ stored by the most efficient upland terrestrial system (e.g. tropical rainforest, Macreadie et al., 2013). Salt marsh ecosystems have thus been globally defined as high-priority zones because of their fragile structure and their important role in providing ecosystem services (Barbier et al., 2011; De Groot et al., 2012). As recently highlighted by

76 the secretariat of the Convention on Biological Diversity, habitat disturbance, erosion and climate
77 change are threatening the functions of salt marshes as C sinks (Secretariat of the Convention on
78 Biological Diversity, 2010). In order to prevent the complete degradation of these ecosystems and to
79 enable implementation of appropriate protection strategies, further information must be obtained
80 about the chemical and biological processes associated with water-soil-plant relationships.

81 Salt marshes are mainly dominated by halophyte vegetation, in which plant zonation is driven by
82 changes in soil morphology, which modify edaphic conditions, competitive mechanisms and thus
83 development of plant communities (Ferronato et al., 2018; Laegdsgaard, 2006; Silvestri et al., 2005).
84 Tidal oscillations, salinity and waterlogging period (hydroperiod) affect plant zonation and form soil
85 hydrosequences ranging from terrestrial to intertidal and subaqueous systems (Ferronato et al., 2018).
86 The distribution of plant communities changes rapidly along hydrosequences and depends on the
87 ability of the plants to tolerate submergence of their roots in salt water (Bertness et al., 1992;
88 Ferronato et al., 2018; Silvestri et al., 2005). The residues of halophyte plant species (mostly annual
89 herbaceous species) contribute greatly to the C sink function of salt marsh ecosystems. Fresh biomass
90 is deposited on the soil surface every year and the rate of degradation depends on the availability of
91 oxygen in soil, persistence of waterlogging and pedoclimatic conditions (Chmura et al., 2003; Mcleod
92 et al., 2011; Sharifi et al., 2013). The tidal oscillations continually supply nutrients and allochthonous
93 suspended mineral matter and organic C, thus contributing to the overall C input in salt marsh soils
94 (Kennedy et al., 2010; Mcleod et al., 2011). The C stable isotope signatures can be used to identify
95 the origin of organic matter, by discriminating terrestrial plant residues (e.g. C3 plants -29 to -22%
96 $\delta^{13}\text{C}$) from riverine phytoplankton (-34 to -26% $\delta^{13}\text{C}$) and marine phytoplankton (-23 to -17%
97 $\delta^{13}\text{C}$) (Bristow et al., 2013; Middelburg and Nieuwenhuize, 1998; Peterson and Fry, 1987). Bristow
98 et al. (2013) recognized the importance of the input of marsh plants and seagrasses such as *Spartina*
99 spp. and *Zostera* spp. on the adjacent salt marshes and mudflats, as well as the presence of microalgae-
100 associated C deposited along the banks, while Santín et al. (2009) highlighted the contribution of the

101 phytoplankton to organic matter (OM) accumulation in a highly exposed area under *Spartina* spp.
102 close to the main tidal channel.

103 Long-standing theory states that soil organic matter comprises a labile pool (e.g. microbial biomass
104 carbon) and a recalcitrant pool (e.g. humic substances) (Ghabbour and Davies, 2001; Rodríguez-
105 Murillo et al., 2017; Schnitzer and Monreal, 2011). In both of these pools, humic substances are
106 commonly defined as complex molecules characterized by the presence of a number of functional
107 oxygenated groups such as carboxyl groups (-COOH) and phenolic hydroxyl groups (Ph-OH), which
108 are involved in soil aggregation, water retention, cation and anion exchange and chelation of mineral
109 elements (Ghabbour and Davies, 2001) and can even act as plant biostimulants (Canellas et al., 2002;
110 Nardi et al., 2002). Many studies have used infrared spectroscopy to analyze the structure of humic
111 compounds, in order to evaluate and predict structural modifications of soil organic matter in space
112 and time (Agnelli et al., 2000; Chai et al., 2007; Liu et al., 2008).

113 There has been some debate about the existence of humic substances in recent years (Lehmann
114 and Kleber, 2015). Thus, soil organic matter formation has been described by these authors "as a
115 continuum of progressively decomposing organic compounds" characterized by a wide range of
116 substances derived from plant residues and particularly enriched in carboxylic acids (Trumbore,
117 1997). Despite the application of advanced spectro-microscopic techniques for *in situ* investigation
118 of soil organic matter, there is no direct evidence for the existence of humic substances in natural
119 soils. Nonetheless, the study of humic substances remains very popular because of the impact of these
120 substances in a wide variety of disciplines (e.g. agronomy, environmental sciences and economy) and
121 the availability of standard samples isolated from soil, peat, leonardite and river water ([http://humic-](http://humic-substances.org/)
122 [substances.org/](http://humic-substances.org/)). In the present study, we chose to maintain the conventional approach to humic and
123 fulvic acids, as recommended by International Humic Substances Society (IHSS), in order to enable
124 comparison of our findings with those of similar studies on soils from hydromorphic environments
125 (Filip et al., 1988; Fookien and Liebezeit, 2003; Santín et al., 2008).

126 Although numerous studies have investigated the nature of humic substances in terrestrial soils,
127 very few have considered the chemical composition of these molecules in marsh and salt marsh
128 sediments (Keller et al., 2009; Santín et al., 2008; Lu and Xu, 2014). In anaerobic microenvironments
129 where oxygen availability is a limiting element, degradation of SOM may also be mediated by
130 reduction reactions. Therefore, the low levels or lack of molecular oxygen in hydromorphic and
131 waterlogged soils may trigger a sequence of changes in chemical properties of the soil that could
132 influence the chemical composition and the structure of humic substances (Rodríguez-Murillo et al.,
133 2011, 2017). Important information can thus probably be obtained about the processes that transform
134 dead plant material into refractory organic C.

135 The chemical diversity of stable SOM depends on the availability of oxygen, which influences the
136 microbial community and the efficiency of soil microorganisms in processing organic carbon (OC)
137 inputs (e.g. by plant cover) (Caravaca et al., 2005; Gleixner et al., 2001). Soil enzymes participate in
138 SOM degradation processes by catalysing the conversion of complex high-molecular weight
139 compounds into smaller compounds, either by redox reactions (catalysed by oxidoreductases), or by
140 the breakdown of complex compounds (catalysed by hydrolases) (Burns, 1982; Sarathchandra et al.,
141 1984). Soil enzyme activity can therefore be used to quantify and monitor changes in soil metabolic
142 activity and SOM dynamics in different ecosystems and in soils disturbed by anthropogenic activities
143 (Trasar-Cepeda et al., 2008). In hydromorphic/subaqueous soils, the restrictive environmental
144 conditions (e.g. low redox potential, soil waterlogging, halinity and sulphide concentration, etc.) can
145 influence the microbial community as well as enzyme synthesis and activities (Arbona et al., 2008;
146 Pulford and Tabatabai, 1988). Soil salinity and waterlogging act as soil dispersant agents, inducing
147 changes in ionic strength and in the molecular stability and sorption of enzymes (Morrissey et al.,
148 2014; Rietl et al., 2016), thereby causing changes in the eco-physiological functions of the microbial
149 community. The pathway of organic matter degradation and the end-products of the microbial
150 metabolism may thus also be affected by both water quality and fluctuations (e.g. tidal oscillations).

151 These changes can be investigated by measuring the enzyme activities associated with
152 biogeochemical cycles (Bello et al., 2015; Burns et al., 2013; Kandeler et al., 1996; Nannipieri, 1994).

153 The aim of the present study was to evaluate changes in SOM and enzyme activities in relation to
154 the soil hydroperiod in three salt marshes in the Baiona Lagoon (Ravenna, North Italy). Soil
155 hydrosequences consisting of terrestrial, intertidal and subaqueous ecosystems have previously been
156 identified in this area, in a study focusing on the relationships between soil hydroperiod, soil
157 physicochemical properties and vegetation pattern (Ferronato et al., 2018). In the present study, the
158 same soils were used to investigate stable and labile carbon pools and also enzyme activities
159 throughout the soil profiles, in order to gain some insight into SOM chemical and biological
160 transformation processes in salt marshes.

161

162 **2. Materials and methods**

163

164 *2.1. Study area and sampling survey*

165

166 The Baiona lagoon (SPA/SAC IT4070004 of the Natura 2000 ecological network) is located in
167 the S. Vitale Regional Park, south of the Po Delta on the northern coast of the Adriatic Sea (N Italy).
168 The area is characterized by retrogradational back-barriers, shoreline and offshore transitional
169 Holocene deposits (Buscaroli et al., 2011; Migani et al., 2015; Veggiani, 1974).

170 Soil profiles were collected in three different salt marshes in the lagoon, according to the terrestrial,
171 intertidal and subaqueous areas defined by Ferronato et al. (2018). In particular, 5 terrestrial (TES),
172 4 intertidal (ITS) and 3 subaqueous (SAS) soil profiles were selected for this study (Figure 1). The
173 soil profiles (0-100 cm depth) were collected using a beaker vibracore sampler equipped with a
174 polyethylene core tube (diameter 6 cm). The genetic horizons were divided and described according
175 to Schoeneberger et al. (2012) and McVey et al. (2012). Subsamples of soil were then prepared in

different ways for subsequent analysis: a) air-dried and sieved at 2 mm, for physical and chemical characterization; b) freeze-dried, for SOM fractionation and diffuse reflectance infrared Fourier transform (DRIFT) analysis; and c) stored at 4 °C, for the analysis of microbial biomass carbon and enzyme activities.

All soil profiles consisted of A-AC-C horizons (Table S1) and were classified according to the US Soil Taxonomy as *Typic psammaquent* to *endaquent* in the TES and ITS systems, and as *Fluventic psammowasssent* to *Typic fluviwassent* in the SAS system (Soil Survey Staff, 2010). All soil horizons (A, AC and C) were analysed separately and the average values for each genetic horizon were subsequently obtained for the TES, ITS and SAS systems.

2.2. Soil physical and chemical analysis

Soil particle size distribution was determined with the pipette method according to Gee and Bauder (1986). Soil pH (pHmeter, Crison, Spain) and electrical conductivity (EC; conductimeter Orion, Germany) were measured in a 1:2.5 (w:v) suspension of soil and deionized water, while total lime (CaCO₃) was quantified by the volumetric method (Loeppert and Suárez, 1996). The total Ca, Na, P and S contents were determined in finely ground soil samples by microwave-assisted acid digestion (Milestone, 1200) with aqua regia (HCl and HNO₃ suprapure, Carlo Erba, Italy, 3:1 v:v) and Inductive Coupled Plasma-Optic Emission Spectroscopy (ICP-OES, Ametek, Germany). The spectrometer was calibrated using international multi element standard (CPI 4400-12627-WG02), and a standard reference material (BCR 320R) was used to check the accuracy of the instrument (< 10%).

Soil total organic carbon (TOC) and total nitrogen (TN) contents were determined by Continuous Flow–Isotope Ratio Mass Spectrometry (CF–IRMS, Delta Plus Thermo Scientific) after pre-treatment of samples with 6 M HCl at 80 °C to eliminate carbonates.

The ¹³C abundance was expressed in delta units (δ¹³C‰):

$$\delta^{13}\text{C}\text{‰} = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 1000$$

202 where, R_{sample} is the isotope ratio $^{13}\text{C}/^{12}\text{C}$ of the sample and R_{standard} is the $^{13}\text{C}/^{12}\text{C}$ ratio of the Pee Dee
203 Belemnite carbonate standard (PDB) (Middelburg and Nieuwenhuize, 1998)

204 All analyses were performed in duplicate for each soil sample and the results shown are the mean
205 values \pm the standard error for the soil samples analysed for each of the soil systems.

206

207 *2.3. SOM fractionation and DRIFT spectroscopy*

208 Humic acid (HA) extraction was performed according to De Nobili et al. (2008). Samples were
209 sequentially extracted at room temperature for 1 h under N₂ flux (1:10 w:v). The HAs were first
210 extracted with 0.5 M NaOH, to yield the fraction considered to consist of labile humic acids (free
211 HA, HAF) consisting of both polysaccharides and humic acids, and then with 0.1 M NaOH + 0.1 M
212 Na₄P₂O₇, to yield the fraction considered to consist of humic acids bound to the soil mineral fraction
213 (bound HA, HAB) via cationic bridges or strongly complexed by polycations (Parsons, 1988). The
214 extracts were precipitated overnight at pH 1.5 with 6 M HCl, and then centrifuged to separate the
215 compounds insoluble at pH 1.5 (humic acids, HA) from those soluble at this pH (fulvic acids and
216 non-humic substances, FA). The HA fractions were then re-suspended in distilled water and freeze-
217 dried.

218 The humic acids (HAF and HAB) were analysed by DRIFT spectroscopy. Spectra were recorded
219 using a Bruker TENSOR series FT-IR Spectrophotometer (Bruker, Ettlingen, Germany) interfaced
220 with a diffuse reflectance apparatus (Spectra-Tech. Inc., Stamford, CT, USA). HA samples were
221 mixed with KBr powder (1:50 w:w) (Aldrich Chemical Co. Milwaukee, WI, USA), which was also
222 used as a background reference. Spectra were recorded as Kubelk-Munk units, ranging from 4000 to
223 400 cm^{-1} and averaged over 100 scans with 4 cm^{-1} resolution and manipulated with OriginLab 7.0
224 software (USA). The percentage area under each band was calculated by curve-fitting analysis,
225 implemented with Grams/386 spectroscopic software (version 6.00, Galactic Industries Corporation,
226 Salem), as described by Mastrolonardo et al. (2015).

The main peaks and bands in the spectra were interpreted according to assignments used by different authors (Fookien and Liebezeit, 2003; Mecozzi and Pierantonio, 2006; Stevenson, 1994) and are shown in Table 1.

2.4. Analysis of microbial biomass carbon and enzyme activities

Soil microbial biomass C (MBC) was determined on previously conditioned (3 days at 25 °C) wet subsamples of soil, by the chloroform fumigation-extraction method (Vance et al., 1987). This method uses chloroform to kill soil microorganisms, before the soil C content is extracted with 0.5 M K₄SO₄. The difference in the C content extracted from fumigated and unfumigated soil samples (measured in a L-TOC analyser, Hypertoc Shimadzu) corresponds to the C released by the dead microorganisms (C flush). This value was then used to estimate the microbial biomass C (expressed in mg kg⁻¹ of dry soil) by applying a transformation factor (Kc) of 0.38 (Joergensen and Brookes, 1990). Microbial biomass C was determined in triplicate for each soil sample, and the results reported are the means ± standard error for the soil samples analysed for each of the soil systems. The microbial quotient (Qmic) was calculated as the percentage of soil MBC relative to the TOC, as reported by Anderson and Domsch (1989). This index was calculated as an indicator of the changes in organic matter due to soil alteration processes and the availability of nutrients to soil microbiota (Sparling, 1997).

Different enzyme assays were performed in triplicate on fresh soil samples by using the traditional methods described below. The results (mean ± standard error of the results for the soil samples analysed for each of the soil systems) are reported as absolute values (μmol of released/consumed compound per gram of oven-dried [105 °C] soil per hour) and as specific activity, i.e. enzyme activity per unit of soil organic carbon (Trasar-Cepeda et al., 2008), which enables comparison of different types of soils under different soil use, and therefore with different organic matter contents (Barriuso et al., 1988). Urease (URE) activity in the samples was determined using 1065.6 mM urea as

253 substrate, incubation for 1.5 h at 37 °C and pH 7.0 (0.2 M phosphate buffer) and measurement of the
254 NH_4^+ released with an ammonia electrode (Nannipieri et al., 1980). Alkaline phosphomonoesterase
255 activity (ALK) was determined using 16 mM *p*-nitrophenyl phosphate as substrate, incubation for 0.5
256 h at 37 °C with Modified Universal Buffer (MUB) pH 11.0 and spectrophotometric measurement (at
257 400 nm) of the *p*-nitrophenol (PNP) released (Tabatabai and Bremner, 1969). β -glucosidase (GLU)
258 activity was determined using *p*-nitrophenyl- β -glucopyranoside 25 mM as substrate, incubation for 1
259 h at 37 °C with MUB pH 6.5 and measurement (at 400 nm) of the *p*-nitrophenol released (Eivazi and
260 Tabatabai, 1988). Arylsulphatase activity (ARYL) was determined using 5 mM *p*-nitrophenyl
261 sulphate as substrate, incubation at pH 5.8 (acetate buffer 0.5 M) and 37 °C for 1 h (Tabatabai and
262 Bremner, 1970). Invertase activity (INV) was determined using 35.06 mM saccharose as substrate,
263 incubation for 3 h at 50 °C and pH 5.5 (acetate buffer 2 M) and measurement of reducing sugars by
264 the Prussian-blue method (Schinner and Mersi, 1990). Catalase activity (CAT) was determined using
265 8.8 mM H_2O_2 as substrate, incubation of the soil samples for 10 min at 20 °C and measurement of the
266 residual H_2O_2 at 505 nm after colorimetric reaction, and the results are expressed as mmol H_2O_2
267 consumed $\text{g}^{-1} \text{h}^{-1}$ (Trasar-Cepeda et al., 1999). In order to obtain an indicator of the enzymatic activity
268 involved in C, N and P cycles, the Synthetic Enzyme Index (SEI) was calculated as the sum of three
269 activities measured using fluorogenic methylumbelliferyl (MUF)-based substrates. β -glucosidase
270 (GLU), chitinase (CHIT) and acid-phosphomonoesterase (ACP) activities were measured using MUF
271 substrates according to the method of Marx et al. (2001). The fluorogenic substrates, prepared with
272 acetate buffer 0.5 M pH 5.5, were 4-MUF- β -D-glucoside 4-MUF-N-acetyl- β -glucosaminide and 4-
273 MUF-phosphate. Fluorescence (excitation 360 nm, emission 450 nm) was measured in an automatic
274 fluorimetric plate-reader (Fluoroskan Ascent) and readings were taken after incubation of plates for
275 0, 30, 60, 120 and 180 minutes at 30 °C. The results are expressed as nmol of MUF released per g of
276 oven-dried soil per hour.

277

278 2.5. Statistical analysis

279

280 Statistical analysis of the data was performed using R free software (R Core Team, 2015), with
281 the Kruskal function of the “*agricolae*” package. As the basic assumptions required for the ANOVA
282 test were not met (i.e. normal data distribution and homogeneity of variance), the data were analysed
283 by one-way nonparametric ANOVA or the Kruskal-Wallis test (Hollander et al., 2013). The latter is
284 a non-parametric method for testing a nominal variable and a continuous (or ordinal) variable by
285 ranking the observations. It tests whether the mean ranks are the same in all the groups, namely if the
286 samples originate from the same distribution. The Kruskal-Wallis test was applied to distinct *a priori*
287 groups formed by the three soil systems (TES, ITS and SAS) and the three different horizons
288 considered (A, AC and C). Holm-Bonferroni correction was applied for multiple comparisons.

289

290 **3. Results**

291

292 *3.1. Physical and chemical properties of salt marsh soils*

293

294 The mean values of the main physical and chemical properties of the diverse horizons in the TES,
295 ITS and SAS soil profiles are shown in Table 2, while Table 3 shows the results of the Kruskal-Wallis
296 test to highlight the significant differences between the three ecosystems (TES, ITS and SAS).

297 The main differences between the physicochemical characteristics of the three different salt marsh
298 systems were observed in the A horizon, in which the pH, EC, organic C and nutrients varied
299 significantly (Table 2). Generally, in all soil pedosequences, the EC decreased sharply from A to AC
300 and C horizons, and the same was observed for clay percentage and the concentrations of
301 macronutrient elements (e.g. TOC, TN, P and S), while Ca and CaCO₃ contents increased in the AC
302 and C horizons (Table 2).

303 Significant differences between the TES, ITS and SAS systems mainly involved the A horizon
304 (Table 3). In the A horizon of the ITS system, the pH was significantly lower than in the A horizons

305 of the other soil systems (Table 3). Moreover, the EC values were significantly higher in all horizons
306 of the ITS system than in all horizons of the TES and SAS systems (Table 3). Similarly, the TOC and
307 P contents were significantly higher in the ITS than in the TES and SAS systems. The Na and S
308 contents increased significantly from the TES to the ITS and SAS system, while there was no
309 significant difference between ITS and SAS systems (Table 3). The $\delta^{13}\text{C}$ signature revealed
310 significant enrichment of the lighter ^{13}C isotope in the surface horizon of TES relative to the same
311 horizon in SAS. Moreover, in the TES and ITS systems, the $\delta^{13}\text{C}$ ratio increased from -25.9 and -25.1
312 to -23.9 and -24.8‰, respectively, throughout the soil profile, while in the SAS system the opposite
313 trend was noted, as the $\delta^{13}\text{C}$ ratio decreased from -23.8 to -25.3‰, although these differences were
314 not statistically significant (Table 3).

315

316 3.2. SOM fractionation and DRIFT spectra

317

318 Humic acids were only extracted from the A horizon because the organic carbon content of the
319 AC and C horizons was too low to obtain accurate results. The DRIFT spectra of both free and bound
320 humic acids (HAF and HAB, respectively) extracted from the A horizon exhibited some
321 modifications along the hydrosequence in both the aliphatic and the aromatic regions (Figure 2).
322 Quantitative information was obtained from the FTIR spectra by Gaussian curve fitting procedure
323 applied to the region considered. The percentage area for each functional group is representative of
324 the structure of humic substances (Figure 3). In general, the peaks in the spectra of the TES system
325 were more variable than those in the spectra of the ITS and SAS systems.

326 In both free and bound HA (HAF and HAB, Figure 3a and 3b, respectively), the percentage area
327 for aliphatic chains ($2954\text{-}2952\text{ cm}^{-1}$) was relatively low in the TES and ITS soils, and was higher in
328 the SAS soils (Figure 3a and b).

329 The peak at around $1729\text{-}1720\text{ cm}^{-1}$, attributed to C=O carbonyl stretching in acids or ketones, was
330 not present in HAF and HAB of the TES soils, while it accounted for 5 and 7 % of the area in the ITS

331 and SAS soils, respectively. The peak at around 1670-1653 cm^{-1} was mainly due to Amide I vibration
332 of proteinaceous material. This band was present in all samples, but it was particularly evident in SAS
333 soils (accounting for 22-24% of the area). Conversely, the peak at 1592-1580 cm^{-1} , assigned to
334 aromatic rings and asymmetric stretching of carboxylate, only appeared in the TES soil and accounted
335 for 19% of the area in HAF (Figure 3).

336 On the other hand, lignin derivatives (indicated by peaks at 1547-1515 cm^{-1}) increased according
337 to the level of soil waterlogging (TES < ITS < SAS). The 1223-1211 cm^{-1} band, attributed to phenol
338 compounds, was only observed in the ITS soils, and accounted for respectively 11.9 and 6.5% of the
339 area in HAF and HAB.

340

341 3.3. Microbial biomass and enzyme activities

342

343 To evaluate the soil microbial community and its activity, microbial biomass C (MBC) content,
344 the metabolic quotient (Q_{mic}) and a synthetic enzyme index (SEI) were determined in all soil profiles
345 (Figure 4 a-c). Enzyme activities and specific enzyme activities were also determined (Table 4 and
346 Figure 5a-f, respectively). The results of the statistical analysis are shown in Table 5.

347 The soil biochemical properties and indexes generally varied widely between the different soil
348 systems (Fig. 4a-c). As found for the soil physical and chemical properties, the most significant
349 differences between the TES, ITS and SAS systems involved the A horizon (Table 5). The MBC
350 content (Figure 4a) was higher in the A horizon of both the TES and ITS than in that of the SAS soil
351 profiles and was much lower in the AC and C horizons. Moreover, in the A horizon, Q_{mic} decreased
352 significantly with increasing hydroperiod (TES>ITS>SAS, Figure 4b). The values of this index were
353 similar in all horizons of the TES soil profiles, but increased throughout the ITS and SAS soil profiles.
354 By contrast, the differences in the SEI between the different horizons and systems were generally not
355 significant (Figure 4c), as a result of the high variability in the data. However, some trends were
356 noted. Thus, for the A horizon, the mean SEI values were lower in the SAS than in TES and ITS soil

357 profiles. Moreover, in the SAS soils, the SEI increased gradually with depth of the soil profile, while
358 in the TES and ITS soils the opposite pattern was observed (Figure 3c). In the AC and C horizons,
359 the MBC and Q_{mic} values were generally lower in the SAS than in the TES and ITS soils, although
360 the differences were statistically significant only for MBC (Table 5). The values of SEI in the AC
361 and C horizons were similar in all three hydrosequences.

362 The values of the enzyme activities are shown in Table 4, and the results of the statistical analysis
363 of the data are reported in Table 5. No significant differences in the enzyme activities were observed
364 between the TES, ITS and SAS systems for the different horizons, with exception of ALK and INV
365 in the A horizons and URE activity in the C horizons. These enzyme activities were significantly
366 lower in the SAS than in TES and ITS soils (Table 5).

367 The enzyme activity generally decreased with depth in the soil profile in the three systems,
368 although the differences were not always statistically significant (data not shown). The CAT, URE,
369 INV and ALK activities were significantly higher in the A horizon than in the deeper horizons of the
370 TES soils. The same trend was observed for CAT, INV and ALK activities in the ITS soil and for
371 CAT and INV in the SAS soils.

372 Generally, the specific enzyme activity, which allows comparison of the enzyme activity between
373 soils with different organic matter content, increased throughout the soil profiles. Notably, in all
374 horizons (A, AC and C), the CAT, URE, GLU and ARYL (Figure 5a, 5b, 5e and 5f) enzyme activities
375 were lower in the ITS soils than in the other soils. Considering the A horizon, CAT/C was lower in
376 the ITS system than in the TES and SAS soils (Figure 5a), although the differences were only
377 statistically significant for the comparison between the ITS and SAS systems (Table 5). Furthermore,
378 INV/C was significantly lower in the SAS than in the TES and ITS (Figure 5d), while GLU/C was
379 significantly higher in the TES system than in the other two systems (Figure 5e).

380 Regarding the AC horizon, CAT/C was significantly lower in the ITS system than both in the TES
381 and SAS systems (Figure 5a), while GLU/C was significantly lower both in the TES and ITS systems
382 than in the SAS system (Figure 5e). This specific activity therefore increased with the intensity of

waterlogging. Considering the C horizon, URE/C, ARYL/C, GLU/C and INV/C were lower in the ITS than in the TES and SAS systems (Figure 5b, 5f, 5e and 5d), and the differences were generally statistically significant, while ALK/C and CAT/C (Figure 5c and 5a) were significantly higher in the SAS system than in the TES and ITS systems, in which the activities were similar (Table 5).

4. Discussion

The soils in the Baiona lagoon area are developed on carbonate sandy substrate, resulting in C horizons with relatively high carbonate content and sandy loam (TES) or sandy clay loam texture (ITS and SAS), in concordance with previous findings (Amorosi et al., 2005; Ferronato et al., 2014; Giambastiani, 2007). The main differences between the three salt marsh systems regarding their organic matter and nutrients content and EC were observed in the A horizon, while the differences between the AC and C horizon of the three salt marsh soils were less pronounced.

The $\delta^{13}\text{C}$ signature indicates the different nature of organic C in the soils considered. In the TES and ITS, the soil surface is enriched by lighter C isotopes ($\delta^{13}\text{C} = -25\text{‰}$ c.a), which are generally associated with organic material derived from terrestrial vascular plants (Bristow et al., 2013). In the deeper horizons, the decrease in C isotopic ratio indicates buried organic C forms and marine plankton material (average value of $\delta^{13}\text{C} = -23\text{‰}$ c.a) (Dickens et al. 2004; Santín et al. 2009). Conversely, the SAS profiles showed the opposite trend, highlighting the presence of C derived from marine plankton on the soil surface and the presence of vascular plants in the deeper layer, probably associated with the salt marsh erosion and subsidence phenomena characteristic of the study area (Ferronato et al, 2016; Teatini et al., 2005).

The difference between the terrestrial and aquatic soil organic matter was determined by analysis of humic substances at the soil surface. Transformation of plant debris is known to be driven by oxygen availability, and in terrestrial systems the organic molecules extracted with alkaline solution

(IHSS, 2015) are typically rich in aromatic compounds and poor in aliphatic ligands (Filip and Alberts, 1994). The structure of humic substances is more complex in terrestrial than in intertidal and subaqueous systems, where degradation of SOM may be mediated by reduction reactions, resulting in humic substances with more aromatic compounds and a higher degree of polymerization than in terrestrial systems (Bronick and Lal, 2005; De Nobili et al., 2008). As the soil waterlogging increased (from ITS to SAS systems), the structure of humic substances became more enriched in aliphatic compounds, and the differences between free HA structure and those bound to the soil mineral fraction (i.e. via cationic bridges or via complexation) became less evident (Figure 2 and 3). Enrichment of aliphatic compounds was mainly observed in the HA extracts from the SAS and can be attributed to the effect of the water movement on the soil surface. Sea water acts as a powerful dispersant of soil aggregates, directly breaking them up and contributing to repulsive charges that disperse clay particles (Bronick and Lal, 2005). The capacity of the soil to retain organic colloids, clay particles and other cementing agents is thus reduced, resulting in enhanced soil fluidity (Surabian, 2007). The shift in some characteristic bands observed in the HA extracts from the SAS profiles (Figure 2) is consistent with the findings of Filip et al. (1988) and Fookien and Liebezeit (2003), who reported that marine humic acids are distinguished by similar bands, mainly attributed to amide linkages of proteins (Ertel and Hedges, 1983). These authors also observed that as the soil hydroperiod increases there is usually an increase in the presence of bands at $1665\text{-}1660\text{ cm}^{-1}$, and at 1725 cm^{-1} . These bands typically increase in the direction fresh plants < dead plants < mud. As expected, the bands at around $1547\text{-}1515\text{ cm}^{-1}$, corresponding to the most characteristic aromatic skeletal vibrations of lignin, increased with the level of soil waterlogging. The persistence of lignin derivatives is probably due to the low level of oxygen, which may decrease the rate of decomposition (Filip et al., 1988). We can conclude that the structure of the organic molecules shifted toward a low complexity along the soil hydrosequences as a consequence of the anaerobic conditions. Nevertheless, we can infer that the water movement (both along the soil hydrosequence and

433 throughout the soil profiles) can produce additional perturbation such as the transport and
434 accumulation of nutrients in different parts of saltmarsh soils.

435 The TOC and MBC contents were lowest in the SAS profiles, while both Q_{mic} and SEI increased
436 significantly with depth of the soil profile, indicating enhancement of biochemical activity in deep
437 horizons. This observation may be related to some pedoturbation due to water movements at the soil
438 surface. Movement of water at the soil surface washes and greatly depletes the soil (Allen, 2000),
439 while in the deeper horizons, water movement is limited and therefore may cause less intense
440 disturbance. This could somehow induce a new equilibrium of microbial communities in AC and C
441 horizons, and a different C metabolic pathways. However, there is another possible explanation for
442 the biochemical characteristics of the SAS. The presence of terrestrial-like C forms in deep SAS
443 horizons (indicated by the isotopic analysis) may be due to past erosion processes, e.g. detachment
444 and deposition of clumps of saltmarsh soils on the sea floor and transportation of fine terrestrial
445 material by water during tidal cycles. The enhanced enzymatic activities detected in the deeper
446 horizons may therefore be associated with past accumulation of upland terrestrial soil material
447 transported by erosion processes. Nevertheless, under altered environmental conditions, the
448 biochemical activity may reach a new equilibrium with different effects in relation to function. This
449 hypothesis suggests that the enhanced enzymatic activity observed in the SAS profiles will be related
450 to the quantity or availability of SOM and also to the type of SOM present in the organo-mineral
451 complex, as under anoxic conditions, the SOM is preserved from degradation and retains immobilized
452 enzymes. The specific enzyme activities (i.e. the activities expressed relative to organic C mass) are
453 often calculated in order to decouple the changes in soil enzyme activities from the changes in organic
454 matter content (Marinari et al., 2012; Trasar-Cepeda et al., 1999; 2008; Vittori Antisari et al., 2011)
455 and to enable comparison of soils under different types of use or subjected to diverse types of
456 disturbance. In terrestrial soils, the activity of hydrolytic enzymes usually decreases with depth
457 (Harrison, 1983; Kuprevich and Shcherbakova, 1971; Trasar-Cepeda and Gil-Sotres, 1987), while
458 the specific enzyme activity generally increases as a result of the lower organic C content in deeper

459 horizons where enzyme activity remains high (Marinari and Vittori Antisari, 2010). This enrichment
460 may represent an ecological mechanism for retaining soil metabolic activity or making organic matter
461 more available to degradation by enzymatic reactions (Boerner et al., 2000; Burns, 1982; Trasar-
462 Cepeda et al., 2008). In this study, the specific enzyme activities generally increased with soil profile
463 depth in all systems (TES, ITS and SAS) and were particularly high in the SAS system (Figure 5).
464 The high specific activity in these soil profiles was particularly noteworthy, as the biochemical
465 activity was expected to be lower in the submerged than in the terrestrial soils. However, as previously
466 hypothesized, the specific enzyme activity in deep horizons of the SAS soils may be due to different
467 types of pedoturbation operated by the combined effects of water movement and soil organisms on
468 the surface horizons, and to the presence of allochthonous terrestrial SOM in deeper horizons. The
469 organic matter derived in the past from terrestrial systems and accumulated in the lagoon sediments
470 may be enriched in immobilized enzymes, which would remain active in AC and C horizons (Burns,
471 1982; Nannipieri, 1994; Nannipieri et al., 1980). Given the lack of reference studies and the
472 complexity of enzyme variability under such environmental conditions, further research is required
473 to confirm this hypothesis.

474 The key focus area between the terrestrial and aquatic environment is the intertidal area, which
475 represents the real transitional system. In this part of the salt marsh hydrosequence, the accumulation
476 of C residues and the occurrence of alternating aerobic/anaerobic soil conditions enable growth of
477 terrestrial plants and mineralization of SOM, as in TES systems. However, the HAs in the ITS soils
478 are relatively poor in aromatic compounds and enriched in phenols and aliphatic compounds
479 (highlighting the lower degree of polymerization than in the TES soils). As in aquatic-like
480 environments, the biochemical and biological processes associated with SOM degradation in the ITS
481 soils are strongly limited by the effect of continuous and provisional waterlogging (Chendrayan et
482 al., 1980). The continuous wet and dry cycles associated with the tidal oscillations contribute greatly
483 to phenomena such as transportation of particulate and soluble organic matter (Fagherazzi et al.,
484 2013) and accumulation of nutrients and marine salts (Ferronato et al., 2016). Accumulation of high

485 amounts of nutrients and organic C content, as well as frequency of flooding and length of
486 waterlogging period, affected both microbial communities and the production and activity of soil
487 hydrolytic enzymes (Geng et al., 2017; Kang and Stanley, 2005), which often decreases in ITS soil
488 horizons. Moreover, the alternation of aerobic/anaerobic conditions due to soil water saturation will
489 affect the metabolic and biochemical pathway of SOM degradation in subaqueous systems, which
490 may therefore differ from those in terrestrial systems, so that a biochemical equilibrium may not be
491 reached.

492

493 **5. Conclusions**

494

495 Soil hydroperiod plays a fundamental role in the degradation of organic matter, inducing a decrease
496 in the oxygen level in soil and thus inhibiting the overall oxidative processes. Water movement also
497 acts as an important pedoturbation agent inducing profound changes in soil physico-chemical and
498 biochemical processes. The biochemical reactions leading to SOM degradation are also influenced
499 by the water flow dynamics, which hinder the activation of enzymes and microbial stabilization. The
500 study findings demonstrate the importance of the origin of the SOM in explaining the enzyme activity
501 in subaqueous soil environments. The high specific enzyme activities in deep horizons of subaqueous
502 environments may be explained by the combined effect of water movement, erosional processes and
503 preservation of SOM under anaerobic conditions. Considering the lack of similar approaches to
504 studying the pathway of SOM degradation in salt marsh ecosystems, further studies are required to
505 test the hypothesis presented in this study at different scales.

506

507 **Acknowledgements**

508

509 The authors thank Professor Gilmo Vianello for his enormous support in helping to advance
510 research on intertidal and subaqueous soil development as a new frontier of the soil science, and for

511 practical field support in this study. The authors also thank Ana I. Iglesias-Tojo for assistance in
512 carrying out the analyses. The authors are grateful for support from the Spanish MINECO through
513 project CGL 2015-66857-C2-1-R, co-financed with FEDER funds from the EU.

514

515 **References**

516 Agnelli, A., Celi, L., Degl’Innocenti, A., Corti, G., Ugolini, F.C., 2000. Chemical and spectroscopic
517 characterization of the humic substances from sandstone-derived rock fragments. *Soil Science*
518 165, 314-327. doi: 10.1097/00010694-200004000-00003.

519 Allen, J.R.L., 2000. Morphodynamics of Holocene salt marshes: a review sketch from the Atlantic
520 and Southern North Sea coasts of Europe. *Quaternary Science Reviews* 19, 1155–1231.
521 doi:10.1016/S0277-3791(99)00034-7.

522 Amorosi, A., Centineo, M.C., Colalongo, M.L., Fiorini, F., 2005. Millennial-scale depositional cycles
523 from the Holocene of the Po Plain, Italy. *Marine Geology* 222–223, 7–18.
524 doi:10.1016/j.margeo.2005.06.041.

525 Anderson, T.-H., Domsch, K.H., 1989. Ratios of microbial biomass carbon to total organic carbon in
526 arable soils. *Soil Biology & Biochemistry* 21, 471–479. doi:10.1016/0038-0717(89)90117-X

527 Arbona, V., Hossain, Z., López-Climent, M.F., Pérez-Clemente, R.M., Gómez-Cadenas, A., 2008.
528 Antioxidant enzymatic activity is linked to waterlogging stress tolerance in citrus. *Physiologia*
529 *Plantarum* 132, 452–466. doi:10.1111/j.1399-3054.2007.01029.x.

530 Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The value
531 of estuarine and coastal ecosystem services. *Ecological Monographs* 81, 169–193.
532 doi:10.1890/10-1510.1.

533 Barriuso, E., Pérez-Mateos, M., González-Carcedo, S., 1988. Actividad ureásica específica del suelo.
534 *Agrochimica* 32, 284-294.

535 Bello, D., Ferronato, C., Vittori Antisari, L., Trasar Cepeda, C., 2015. Preliminary results on
536 enzymatic activities in two salt marsh soils differing in hydromorphic characteristics and

537 vegetation cover. *EQA – International Journal of Environmental Quality* 19, 1–15.
538 doi:10.6092/issn.2281-4485/5801.

539 Bertness, M.D., Gough, L., Shumway, S.W., 1992. Salt tolerances and the distribution of fugitive salt
540 marsh plants. *Ecology* 73, 1842–1851.

541 Boerner, R.E.J., Decker, K.L.M., Sutherland, E.K., 2000. Prescribed burning effects on soil enzyme
542 activity in a southern Ohio hardwood forest: a landscape-scale analysis. *Soil Biology &*
543 *Biochemistry* 32, 899–908. doi:10.1016/S0038-0717(99)00208-4.

544 Bristow, L.A., Jickells, T.D., Weston, K., Marca-Bell, A., Parker, R., Andrews, J.E., 2013. Tracing
545 estuarine organic matter sources into the southern North Sea using C and N isotopic signatures.
546 *Biogeochemistry* 113, 9–22. doi:10.1007/s10533-012-9758-4.

547 Bronick, C.J., Lal, R., 2005. Soil structure and management: A review. *Geoderma* 124, 3–22.
548 doi:10.1016/j.geoderma.2004.03.005.

549 Burns, R.G., 1982. Enzyme activity in soil: Location and a possible role in microbial ecology. *Soil*
550 *Biology & Biochemistry* 14, 423–427. doi:10.1016/0038-0717(82)90099-2.

551 Burns, R.G., DeForeset, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein, M.D.,
552 Weintraub, M.N., Zoppini, A., 2013. Soil enzymes in a changing environment: Current
553 knowledge and future directions. *Soil Biology & Biochemistry* 58, 216–234.
554 doi:10.1016/J.SOILBIO.2012.11.009.

555 Buscaroli, A., Dinelli, E., Zannoni, D., 2011. Geohydrological and environmental evolution of the
556 area included among the lower course of the Lamone river and the Adriatic coast. *EQA –*
557 *International Journal of Environmental Quality* 5, 11–22.

558 Canellas L.P., Lopes Olivares F., Okorokova-Façanha, A.L., Rocha Façanha, A., 2002. Humic Acids
559 isolated from Earthworm composte enhance Root Elongation, Lateral Emergence, and Plasma
560 membrane H⁺-ATPase Activity in Maize Roots. *Plant Physiology* 130, 1951-1957.
561 doi.org/10.1104/pp.007088.

562 Caravaca, F., Alguacil, M.M., Torres, P., Roldán, A., 2005. Plant type mediates rhizospheric

563 microbial activities and soil aggregation in a semiarid Mediterranean salt marsh. *Geoderma* 124,
564 375–382. doi:10.1016/j.geoderma.2004.05.010.

565 Chai, X., Takayuki, S., Cao, X., Guo, Q., Zhao, Y., 2007. Spectroscopic studies of the progress of
566 humification processes in humic substances extracted from refuse in a landfill. *Chemosphere* 69,
567 1446–1453. doi:10.1016/j.chemosphere.2007.04.076.

568 Chendrayan K., Adhya T. K., Sethunathan N. (1980) Dehydrogenase and invertase activities of
569 flooded soils. *Soil Biology & Biochemistry* 12, 211-213.

570 Chmura, G.L., Anisfeld, S.C., Cahoon, D.R., Lynch, J.C., 2003. Global carbon sequestration in tidal,
571 saline wetland soils. *Global Biogeochemical Cycles* 17, 1111. doi:10.1029/2002GB001917.

572 De Groot, R., Brander, L., van der Ploeg, S., Costanza, R., Bernard, F., Braat, L., Christie, M.,
573 Crossman, N., Ghermandi, A., Hein, L., Hussain, S., Kumar, P., McVittie, A., Portela, R.,
574 Rodriguez, L.C., ten Brink, P., van Beukering, P., 2012. Global estimates of the value of
575 ecosystems and their services in monetary units. *Ecosystem Services* 1, 50–61.
576 doi:10.1016/j.ecoser.2012.07.005.

577 De Nobili, M., Contin, M., Mahieu, N., Randall, E.W., Brookes, P.C., 2008. Assessment of chemical
578 and biochemical stabilization of organic C in soils from the long-term experiments at
579 Rothamsted (UK). *Waste Management* (New York, N.Y.) 28, 723–33.
580 doi:10.1016/j.wasman.2007.09.025.

581 Dickens, A.F., Gélinas, Y., Masiello, C.A., Wakeham, S., Hedges, J.I., 2004. Reburial of fossil
582 organic carbon in marine sediments. *Nature* 427, 336–339. doi:10.1038/nature02299.

583 Eivazi, F., Tabatabai, M.A., 1988. Glucosidases and galactosidases in soils. *Soil Biology &*
584 *Biochemistry* 20, 601–606. doi:10.1016/0038-0717(88)90141-1.

585 Ertel, J.R., Hedges, J.I., 1983. Bulk chemical and spectroscopic properties of marine and terrestrial
586 humic acids, melanoidins and catechol-based synthetic polymers, in: Christman, R.F., Gjessing,
587 E.T. (Eds.), *Aquatic and Terrestrial Humic Materials*, Ann Arbor Science, Ann Arbor, MI,
588 pp.143-163

589 Fagherazzi, S., Wiberg, P.L., Temmerman, S., Struyf, E., Zhao, Y., Raymond, P.A., 2013. Fluxes of
590 water, sediments, and biogeochemical compounds in salt marshes. *Ecological Processes* 2, 3.
591 doi:10.1186/2192-1709-2-3.

592 Ferronato, C., Falsone, G., Natale, M., Zannoni, D., Buscaroli, A., Vianello, G., Vittori, L., 2016.
593 Chemical and pedological features of subaqueous and hydromorphic soils along a
594 hydrosequence within a coastal system (San Vitale Park, Northern Italy). *Geoderma* 265, 141–
595 151. doi:10.1016/j.geoderma.2015.11.018.

596 Ferronato, C., Speranza, M., Ferroni, L., Buscaroli, A., Vianello, G., Vittori Antisari, L., 2018.
597 Vegetation response to soil salinity and waterlogging in three saltmarsh hydrosequences through
598 macronutrients distribution. *Estuarine, Coastal and Shelf Science* 200, 131–140.
599 doi:10.1016/j.ecss.2017.10.019.

600 Ferronato, C., Vianello, G., Vittori Antisari, L., 2014. The evolution of the Po Valley and Reno basin
601 (North Italy) through the historical cartography: vicissitude of a land reclamation, in: *Regional*
602 *Symposium on Water, Wastewater and Environment: Traditions and Culture*. pp. 741–752.

603 Filip, Z., Alberts, J.J., 1994. Microbial utilization resulting in early diagenesis of salt-marsh humic
604 acids. *Science of The Total Environment* 144, 121–135. doi:10.1016/0048-9697(94)90433-2.

605 Filip, Z., Alberts, J.J., Cheshire, M. V, Goodman, B.A., Bacon, J.R., 1988. Comparison of salt marsh
606 humic acid with humic-like substances from the indigenous plant species *Spartina alterniflora*
607 (Loisel). *Science of The Total Environment* 71, 157–172. doi:10.1016/0048-9697(88)90164-7.

608 Fookien, U., Liebezeit, G., 2003. An IR Study of Humic Acids Isolated from Sediments and Soils.
609 *Senckenbergiana Maritima* 32, 183–189. doi:10.1007/BF03043094.

610 Gee, G. W., Bauder, J. W., 1986. *Methods of Soil Analysis: Part 1—Physical and Mineralogical*
611 *Methods*, SSSA Book Series. Soil Science Society of America, American Society of Agronomy,
612 Madison, WI. doi:10.2136/sssabookser5.1.2ed.c15.

613 Geng, Y., Wang, D., Yang, W., 2017. Effects of different inundation periods on soil enzyme activity
614 in riparian zones in Lijiang. *Catena* 149, 19–27. doi:10.1016/j.catena.2016.08.004.

615 Ghabbour, E.A., Davies, G., 2001. Humic substances: structures, models and functions. Royal
616 Society of Chemistry.

617 Giambastiani, B., 2007. Evoluzione idrologica ed idrogeologica della pineta di S. Vitale (Ravenna),
618 PhD Thesis, University of Bologna.

619 Gleixner, G., Czimczik, C.J., Kramer, C., Lühker, B., Schmidt, M.W.I., 2001. Plant Compounds and
620 Their Turnover and Stabilization as Soil Organic Matter, in: Schulze, E.D., Heimann, M.,
621 Harrison, S., Holland, E., Lloyd, J., Prentice, I.C., Schimel, D. (Eds.), Global Biogeochemical
622 Cycles and the Physical Climate System, Academic Press, San Diego, CA, pp. 201–215.

623 Harrison, A.F., 1983. Relationship between intensity of phosphatase activity and physico-chemical
624 properties in woodland soils. *Soil Biology & Biochemistry* 15, 93–99. doi:10.1016/0038-
625 0717(83)90124-4.

626 Hollander, M., Wolfe, D.A., Chicken, E., 2013. Nonparametric statistical methods. John Wiley &
627 Sons.

628 IHSS, 2015. What are Humic Substances? [WWW Document]. URL <http://humic-substances.org/>
629 (accessed 12.5.17).

630 Joergensen, R.G., Brookes, P.C., 1990. Ninhydrin-reactive nitrogen measurements of microbial
631 biomass in 0.5 M K₂SO₄ soil extracts. *Soil Biology & Biochemistry* 22, 1023–1027.
632 doi:10.1016/0038-0717(90)90027-W.

633 Kandeler, F., Kampichler, C., Horak, O., 1996. Influence of heavy metals on the functional diversity
634 of soil microbial communities. *Biology and Fertility of Soils* 23, 299–306.
635 doi:10.1007/BF00335958.

636 Kang, H., Stanley, E.H., 2005. Effects of levees on soil microbial activity in a large river floodplain.
637 *River Research and Applications* 21, 19–25. doi:10.1002/rra.811.

638 Kelleher, B.P., Simpson, A.J., 2006. Humic Substances in Soils: Are They Really Chemically
639 Distinct? *Environmental Science and Technology* 40, 4605–4611. doi:10.1021/ES0608085.

640 Keller, J.K., Weisenborn, P.B., Megonigal, J.P., 2009. Humic acids as electron acceptors in wetland

decomposition. *Soil Biology & Biochemistry* 41, 1518–1522. doi:10.1016/j.soilbio.2009.04.008

Kennedy, H., Beggins, J., Duarte, C.M., Fourqurean, J.W., Holmer, M., Marbá, N., Middelburg, J.J.,
2010. Seagrass sediments as a global carbon sink: Isotopic constraints. *Global Biogeochemical
Cycles* 24, 1–9. doi:10.1029/2010GB003848.

Kuprevich, V.F., Shcherbakova, T.A., 1971. Comparative enzymatic activity in diverse types of soil,
in: McLaren A.D., Skujins J.J. (Eds), *Soil Biochemistry*, vol. 2, Marcel Dekker, New York, pp.
167–201.

Laegdsgaard, P., 2006. Ecology, disturbance and restoration of coastal saltmarsh in Australia: a
review. *Wetlands Ecology and Management* 14, 379–399. doi:10.1007/s11273-005-8827-z.

Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature* 528, 60.
doi:10.1038/nature16069.

Lehmann, J., Solomon, D., Kinyangi, J., Dathe, L., Wirick, S., Jacobsen, C., 2008. Spatial complexity
of soil organic matter forms at nanometre scales. *Nature Geoscience* 1, 238–242.
doi:10.1038/ngeo155.

Liu, Q., Shurong, W., Yun, Z., Zhongyang, L., Kefa, C., 2008. Mechanism study of wood lignin
pyrolysis by using TG–FTIR analysis. *Journal of Analytical and Applied Pyrolysis* 82, 170–177.
doi:10.1016/J.JAAP.2008.03.007.

Loeppert, R.H., Suarez, D.L., 1996. Carbonate and Gypsum, in: D. L. Sparks, A. L. Page, P. A.
Helmke, R. H. Loeppert, P. N. Soltanpour, M. A. Tabatabai, C. T. Johnston, M. E. Sumner
(Eds.), *Methods of Soil Analysis. Part 3. Chemical Methods*. SSSA Book Series. Soil Science
Society of America-American Society of Agronomy, Madison, WI, pp.437-474.

Lu, Y., Xu, H., 2014. Distribution characteristic of soil organic carbon fraction in different types of
wetland in Hongze Lake of China. *The Scientific World Journal* 2014, 1-5.
doi:10.1155/2014/487961

Macreadie, P.I., Hughes, A.R., Kimbro, D.L., D’Odorico, P., McGlathery, K., 2013. Loss of “Blue
Carbon” from Coastal Salt Marshes Following Habitat Disturbance. *PLoS ONE* 8, e69244.

doi:10.1371/journal.pone.0069244

Marinari, S., Carbone, S., Vittori Antisari, L., Grego, S., Vianello, G., 2012. Microbial activity and functional diversity in Psamment soils in a forested coastal dune-swale system. *Geoderma* 173–174, 249–257. doi:10.1016/j.geoderma.2011.12.023

Marinari, S., Vittori Antisari, L., 2010. Effect of lithological substrate on microbial biomass and enzyme activity in brown soil profiles in the northern Apennines (Italy). *Pedobiologia* 53, 313–320. doi:10.1016/j.pedobi.2010.02.004

Marx, M.C., Wood, M., Jarvis, S.C., 2001. A microplate fluorimetric assay for the study of enzyme diversity in soils. *Soil Biology & Biochemistry* 33, 1633-1640. doi.org/10.1016/S0038-0717(01)00079-7.

Mastrolonardo, G., Francioso, O., Di Foggia, M., Bonora, S., Forte, C., Certini, G., 2015. Soil pyrogenic organic matter characterisation by spectroscopic analysis: a study on combustion and pyrolysis residues. *Journal of Soils and Sediments* 15, 769–780. doi:10.1007/s11368-014-1034-x

McLeod, E., Chmura, G.L., Bouillon, S., Salm, R., Björk, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., Silliman, B.R., 2011. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Frontiers in Ecology and the Environment* 9, 552–560. doi:10.1890/110004

McVey, S., Schoeneberger, P.J., Turenne, J., Payne, M., Wysocki, D.A., 2012. Subaqueous soils (SAS) description, in: *Field Book for Describing and Sampling Soils*. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE., USA, pp. 2–97.

Mecozzi, M., Pierantonio, E., 2006. Carbohydrates proteins and lipids in fulvic and humic acids of sediments and its relationships with mucilaginous aggregates in the Italian seas. *Marine Chemistry* 101, 27–39. doi:10.1016/j.marchem.2006.01.001

Middelburg, J.J., Nieuwenhuize, J., 1998. Carbon and nitrogen stable isotopes in suspended matter and sediments from the Schelde Estuary. *Marine Chemistry* 60, 217–225. doi:10.1016/S0304-

693 4203(97)00104-7

694 Migani, F., Borghesi, F., Dinelli, E., 2015. Geochemical characterization of surface sediments from
695 the northern Adriatic wetlands around the Po river delta. Part I: Bulk composition and relation
696 to local background. *Journal of Geochemical Exploration* 156, 72–88.
697 doi:10.1016/j.gexplo.2015.05.003

698 Mitra, S., Wassmann, R., Vlek, P.L.G., 2005. An appraisal of global wetlands area and its organic
699 carbon stock. *Current Science* 25–35.

700 Morrissey, E.M., Gillespie, J.L., Morina, J.C., Franklin, R.B., 2014. Salinity affects microbial activity
701 and soil organic matter content in tidal wetlands. *Global Change Biology* 20, 1351–62.
702 doi:10.1111/gcb.12431

703 Nannipieri, P., 1994. The potential use of soil enzymes as indicators of productivity, sustainability
704 and pollution, in: Pankhurst, C.E., Double, B.M., Gupta, V.V.S.R., Grace, P.R. (Eds.), *Soil*
705 *Biota: Management and Sustainable Farming Systems*, CSIRO, Adelaide, pp. 238–244.

706 Nannipieri, P., Ceccanti, B., Cervelli, S., Matarese, E., 1980. Extraction of Phosphatase, Urease,
707 Proteases, Organic Carbon, and Nitrogen from Soil. *Soil Science Society of America Journal*
708 44, 1011. doi:10.2136/sssaj1980.03615995004400050028x.

709 Nardi, S., Pizzeghello, D., Muscolo, A., Vianello, A., 2002. Physiological effects of humic substances
710 on higher plants. *Soil biology & Biochemistry* 34, 1527-1536. doi.org/10.1016/S0038-
711 0717(02)00174-8.

712 Parsons, J.R., 1988. Isolation of humic substances from soils and sediments, in: Frimmel F.H.,
713 Christman, R.F. (Eds), *Humic Substances and Their Role in the Environment*, John Wiley &
714 Sons Ltd., Chichester, pp. 3-14.

715 Peterson, B.J., Fry, B., 1987. Stable Isotopes in Ecosystem Studies. *Annual Review of Ecology and*
716 *Systematics* 18, 293–320. doi:10.1146/annurev.es.18.110187.001453

717 Pulford, I.D., Tabatabai, M.A., 1988. Effect of waterlogging on enzyme activities in soils. *Soil*
718 *Biology & Biochemistry* 20, 215–219. doi:10.1016/0038-0717(88)90039-9

719 R Core Team, 2015. R: A language and environment for statistical computing. R Foundation for
720 Statistical Computing [WWW Document]. URL <https://www.r-project.org/>

721 Rietl, A.J., Overlander, M.E., Nyman, A.J., Jackson, C.R., 2016. Microbial Community Composition
722 and Extracellular Enzyme Activities Associated with *Juncus roemerianus* and *Spartina*
723 *alterniflora* Vegetated Sediments in Louisiana Saltmarshes. *Microbial Ecology* 71, 290–303.
724 doi:10.1007/s00248-015-0651-2

725 Rodríguez-Murillo, J.C., Almedrons, G., Knicker, H., 2011. Wetland soil organic matter composition
726 in a Mediterranean semiarid wetland (Las Tablas de Daimiel, Central Spain): Insight into
727 different carbon sequestration pathways. *Organic Geochemistry* 42, 762–773.
728 doi:10.1016/J.ORGGEOCHEM.2011.05.007

729 Rodríguez-Murillo, J.C., Almendros, G., Knicker, H., 2017. Humic acid composition and
730 humification processes in wetland soils of a Mediterranean semiarid wetland. *Journal of Soils*
731 *and Sediments* 17, 2104–2115. doi:10.1007/s11368-017-1663-y

732 Santín, C., González-Pérez, M., Otero, X.L., Ángel Álvarez, M., Macías, F., 2009. Humic substances
733 in estuarine soils colonized by *Spartina maritima*. *Estuarine, Coastal and Shelf Science* 81, 481–
734 490. doi:10.1016/j.ecss.2008.12.013

735 Santín, C., González-Pérez, M., Otero, X.L., Vidal-Torrado, P., Macías, F., Álvarez, M.Á., 2008.
736 Characterization of humic substances in salt marsh soils under sea rush (*Juncus maritimus*).
737 *Estuarine, Coastal and Shelf Science* 79, 541–548. doi:10.1016/j.ecss.2008.05.007

738 Sarathchandra, S.U., Perrott, K.W., Upsdell, M.P., 1984. Microbiological and biochemical
739 characteristics of a range of New Zealand soils under established pasture. *Soil Biology &*
740 *Biochemistry* 16, 177–183. doi:10.1016/0038-0717(84)90109-3

741 Schinner, F., Mersi, W. von, 1990. Xylanase-, CM-Cellulase- and Invertase Activity in Soil: An
742 Improved Method. *Soil Biology & Biochemistry*. 22, 511-515.

743 Schnitzer, M., Monreal, C.M., 2011. Chapter three – Quo vadis Soil Organic Matter Research?: A
744 Biological Link to the Chemistry of Humification. *Advances in Agronomy* 113, 139-213.

doi:10.1016/B978-0-12-386473-4.00008-7

Schoeneberger, P., Wysocki, D.A., Benham, E.C.J., 2012. Field book for describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE., USA.

Secretariat of the Convention on Biological Diversity, 2010. Global Biodiversity Outlook 3. Secretariat of the Convention on Biological Diversity, Montréal.

Sharifi, A., Kalin, L., Hantush, M.M., Isik, S., Jordan, T.E., 2013. Carbon dynamics and export from flooded wetlands: A modeling approach. *Ecological Modelling* 263, 196–210. doi:10.1016/j.ecolmodel.2013.04.023

Silvestri, S., Defina, A., Marani, M., 2005. Tidal regime, salinity and salt marsh plant zonation. *Estuarine, Coastal and Shelf Science* 62, 119–130. doi:10.1016/j.ecss.2004.08.010

Soil Survey Staff, 2010. Keys to Soil Taxonomy, 12th ed. United States Department of Agriculture, Natural Resources Conservation Service, Lincoln, NE., USA.

Sparling, G.P., 1997. Soil microbial biomass, activity and nutrient cycling as indicators of soil health.

Stevenson, F.J., 1994. Humus chemistry : genesis, composition, reactions. Wiley and Sons, New York.

Surabian, D.A., 2007. Moorings : An Interpretation from the Coastal Zone Soil Survey of Little Narragansett Bay , Connecticut and Rhode Island 92, 90–92.

Tabatabai, M.A., Bremner, J.M., 1970. Arylsulfatase Activity of Soils. *Soil Science Society of America Journal* 34, 225. doi:10.2136/sssaj1970.03615995003400020016x

Tabatabai, M.A., Bremner, J.M., 1969. Use of *p*-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biology & Biochemistry* 1, 301–307. doi:10.1016/0038-0717(69)90012-1

Teatini, P., Ferronato, M., Gambolati, G., Bertoni, W., Gonella, M., 2005. A century of land subsidence in Ravenna, Italy. *Environmental Geology* 47, 831–846. doi:10.1007/s00254-004-1215-9

Trasar-Cepeda, C., Camiña, F., Leirós, M.C., Gil-Sotres, F., 1999. An improved method to measure

771 catalase activity in soils. *Soil Biology & Biochemistry* 31, 483–485. doi:10.1016/S0038-
772 0717(98)00153-9

773 Trasar-Cepeda, C., Leirós, M.C., Gil-Sotres, F., 2008. Hydrolytic enzyme activities in agricultural
774 and forest soils. Some implications for their use as indicators of soil quality. *Soil Biology &*
775 *Biochemistry* 40, 2146–2155. doi:10.1016/j.soilbio.2008.03.015

776 Trasar-Cepeda, M.C., Gil-Sotres, F., 1987. Phosphatase activity in acid high organic matter soils in
777 Galicia (NW Spain). *Soil Biology & Biochemistry* 19, 281–287. doi:10.1016/0038-
778 0717(87)90010-1

779 Trumbore, S.E., 1997. Potential responses of soil organic carbon to global environmental change.
780 *Proceedings of the National Academy of Sciences of the United States of America* 94, 8284–91.
781 doi:10.1073/PNAS.94.16.8284

782 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil
783 microbial biomass C. *Soil Biology & Biochemistry* 19, 703–707. doi:10.1016/0038-
784 0717(87)90052-6

785 Veggiani, A., 1974. Le ultime vicende geologiche del ravennate, in: *Influenza Di Insediamenti*
786 *Industriali Sul Circostante Ambiente Naturale – Studio Sulla Pineta Di San Vitale Di Ravenna.*
787 Ed Compositori, Bologna.

788 Vittori Antisari, L., Marinari, S., Dell’Abate, M.T., Baffi, C., Vianello, G., 2011. Plant cover and
789 epipedon SOM stability as factors affecting brown soil profile development and microbial
790 activity. *Geoderma* 161, 212–224. doi:10.1016/j.geoderma.2010.12.021.

791

792

793

794

795

796

797 **Figure captions:**

798 Figure 1: Study area, sampling sites and WGS-84 UTM 33T coordinates.

799 Figure 2. DRIFT spectra of free (F) and bound (B) humic acids (according to conventional theory)
800 extracted from the A horizons of the TES, ITS and SAS systems.

801 Figure 3. Percentage of the main peak areas on a) free HA and b) bound HA (according to
802 conventional theory) extracted from the TES, ITS and SAS soils.



803 Figure 4. Mean values \pm standard error (SE) of microbial biomass carbon (MBC), microbial quotient
804 (Q_{mic}) and Synthetic Enzyme Index (SEI) for A, AC and C horizons of the TES, ITS and SAS
805 profiles.

806 Figure 5. Mean values \pm standard error (SE) of specific enzyme activities for A, AC and C horizons
807 of the TES, ITS and SAS soil profiles. Data are expressed as mmol H_2O_2 consumed $g\ C^{-1}\ h^{-1}$ (CAT/C),
808 $\mu mol\ NH_3\ g\ C^{-1}\ h^{-1}$ (URE/C), $\mu mol\ PNP\ g\ C^{-1}\ h^{-1}$ (ALK/C, ARYL/C, GLU/C), $\mu mol\ glucose\ g^{-1}\ h^{-1}$
809 (INV/C).


810



811




812



WGS84-UTM33					
TES			ITS		
SITE	E (m)	N (m)	SITE	E (m)	N (m)
TES-1A	282468	4932596	ITS-1A	282478	4932594
TES-2A	282473	4932595	ITS-2B	281264	4930269
TES-3B	281276	4930301	ITS-3C	282411	4930926
TES-4B	281269	4930281	ITS-4C	282412	4930921
TES-5C	282412	4930932			
			SAS		
SITE	E (m)	N (m)	SITE	E (m)	N (m)
SAS-1A	282487	4932592			
SAS-2B	281260	4930261			
SAS-3C	282412	4930913			



3a. HAF**3b. HAB**











Table 1. Assignment of the main bands observed in the SOM FT-IR spectra according to Fookien and Liebezeit (2003), Mecozzi and Pierantonio (2006), and Stevenson (1994)

Wavelength (cm ⁻¹)	Band assignments
3500-3200	O–H stretching vibration in phenols, carboxylic acids and carbohydrates
2952-2850	Symmetric and symmetric C–H stretching
1730-1720	C=O stretching in acids or ketones
1670-1650	C=O stretching in Amide I; aromatic and aliphatic C=C stretching; C=N stretching; H-bonded conjugated ketones, carboxyls and quinones
1570-1515	Aromatic C=C stretching, aromatic skeletal vibrations, aromatic skeletal (lignin), amide II; carboxylates
1440-1340	C–H deformation of CH ₂ and CH ₃ groups and/or to symmetric stretching of COO– groups
1260-1216	Amide III, C–O stretching and O–H deformation of COOH groups and to C–O stretching of aryl ethers and phenols

Table 2. Physical and chemical properties of soil horizons (A, AC and C horizons) in the different ecosystems (TES, ITS, SAS). Mean values and Standard Error (SE) for each group are shown. “na”= not available (because of lack of sufficient material).

		pH		EC dS m ⁻¹		Sand %		Silt %		Clay %		CaCO ₃ g kg ⁻¹		Ca g kg ⁻¹	
		Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
A	TES	8	0.1	12	3	58.2	8.8	30.1	6.8	11.7	3.1	6.3	1.2	31.5	3.6
	ITS	7.5	0.2	24.9	2.6	16.5	1.9	64.2	5.2	19.3	3.6	3.3	1.4	17.2	4.1
	SAS	8.3	0.2	8.1	1.4	na	na	na	na	na	na	na	na	36.7	13.2
AC	TES	8.4	0.1	6.1	0.6	57.3	11.8	36.3	10.6	6.4	1.3	12.9	0.8	51.4	1.8
	ITS	8.2	0.1	10.2	0.9	40.8	14.6	51.6	14.3	7.6	1	9.6	2.3	36	7.8
	SAS	8.3	0.1	7.2	0.8	39.3	15.9	53.6	14.4	7.1	1.5	15.1	1	57.1	2.2
C	TES	8.5	0.1	5.7	0.9	65.5	13	28.2	12.5	6.3	1.1	15.1	0.9	59.6	2.7
	ITS	8.5	0	8.1	1.5	53.7	15.9	42	15.8	4.4	0.7	13.4	0.7	52.8	0.9
	SAS	8.5	0.1	5.1	0.8	72.6	6.6	26.5	9.8	5.2	1.3	15.7	0.6	59.8	1.7

		TOC g kg ⁻¹		δ ¹³ C ‰		TN g kg ⁻¹		Na g kg ⁻¹		P g kg ⁻¹		S g kg ⁻¹	
		Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
A	TES	51.6	15.1	-25.9	0.3	4.7	1.5	6.5	0.9	0.7	0.1	1.3	0.6
	ITS	111.1	17.4	-25.1	0.6	11.1	1.9	9.9	1	1	0.1	4.6	1.2
	SAS	38.3	17.7	-23.8	0.9	6	3.1	13.9	1.3	0.5	0.1	8.2	1.2
AC	TES	5.9	0.9	-25	0.2	0.6	0.1	5.6	0.7	0.5	0.1	0.4	0.1
	ITS	6.1	1.4	-25	0.3	0.7	0.2	8.6	1.3	0.6	0.1	0.6	0.1
	SAS	5.1	1	-25.3	0.3	0.9	0.3	6.7	1	0.5	0.1	1.7	0.5
C	TES	3.8	1.1	-24.8	0.1	0.4	0.1	5.2	0.9	0.5	0.1	0.3	0.1
	ITS	4.7	0.7	-23.9	0.2	0.5	0.1	5.3	0.7	0.4	0.1	0.3	0.1
	SAS	2	0.4	-25.3	0.3	0.3	0	4.3	0.6	0.4	0	0.7	0.1

Table 3. Results of the Kruskal-Wallis test based on the grouping of A, AC and C horizons from the three salt marsh ecosystems (TE, IT and SA).

nd= not determined; ns= not significant; * $P < 0.05$, ** $P < 0.01$.

[illegible]

Table 4. Mean values (\pm standard error, SE) of enzyme activities in the A, AC and C horizons of the TE, IT and SA soil profiles.

		CAT		URE		ALK P		GLU		ARYL		INV	
		mmol H ₂ O ₂ g ⁻¹ h ⁻¹		μ mol NH ₃ g ⁻¹ h ⁻¹		μ mol PNP g ⁻¹ h ⁻¹		μ mol PNP g ⁻¹ h ⁻¹		μ mol PNP g ⁻¹ h ⁻¹		μ mol glucose g ⁻¹ h ⁻¹	
		mean	SE	mean	SE	mean	SE	mean	SE	mean	SE	mean	SE
A	TES	17.4	4.2	2.9	0.7	3	0.4	0.6	0.2	0.4	0.1	12.1	4.4
	ITS	22.5	4.8	2.2	0.6	3.3	1	0.4	0.1	0.6	0.2	11.6	2.2
	SAS	13.9	0.7	1	0.3	0.3	0.1	0.3	0.1	0.6	0	1.6	0.1
AC	TES	7	2.8	0.5	0.1	0.7	0.2	0.3	0.1	0.5	0.1	0.7	0.2
	ITS	4.5	1.3	0.6	0.3	0.9	0.3	0.3	0.1	0.5	0.1	1.4	0.5
	SAS	10.1	2.5	0.4	0.1	0.5	0	0.6	0.1	0.5	0.1	0.5	0.1
C	TES	4.9	2.3	0.6	0.1	0.3	0	0.6	0.1	0.5	0.2	0.3	0.1
	ITS	6.7	5.3	0.4	0.1	0.6	0.2	0.3	0.1	0.4	0.1	0.4	0
	SAS	5.1	1.1	0.3	0	0.3	0.1	0.6	0.1	0.6	0.1	0.2	0

Table 5. Results of the Kruskal Wallis test based on the single A, AC and C horizons of three salt marsh ecosystems (TE, IT and SA). ns= not significant; * P <0.05, ** P <0.01, *** P <0.001.

		MBC	Qmic	SEI	CAT	URE	ALK	INV	GLU	ARYL	CAT/C	URE/C	ALK/C	INV/C	GLU/C	ARYL/C
A	TES vs ITS	ns	***	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns
	TES vs SAS	***	***	ns	ns	ns	*	ns	ns	ns	ns	ns	ns	***	**	ns
	ITS vs SAS	***	*	ns	ns	ns	*	ns	ns	ns	*	ns	ns	*	ns	ns
AC	TES vs ITS	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns
	TES vs SAS	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns
	ITS vs SAS	*	ns	ns	ns	ns	ns	ns	ns	ns	*	ns	ns	ns	ns	ns
C	TES vs ITS	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	*	ns	ns	ns	ns
	TES vs SAS	*	ns	ns	ns	**	ns	ns	ns	ns	*	ns	ns	*	ns	ns
	ITS vs SAS	*	ns	ns	ns	**	ns	*	ns	ns	**	*	ns	ns	ns	ns

Table S1. Morphological soil classification (McVey et al., 2012; Schoeneberger et al., 2012).

Soil Classification	Soil System	Master	Depth (cm)	Bound.	Color Munsell		Text.	HC (cm h ⁻¹)	Structure	Consistence /Fluidity	Concentrations					Sulphide detection				
					Moist	Dry					Mottles		Roots	Bio	Other	RXF	H ₂ O ₂	Odor	pH _m	pH _{fin}
											I	II								
Typic Psammaquent	TES	A1	0-5	A/W	10YR 3/2	10YR 4/3	SL	7.59	SBK/1/f	so/po			c/f-m				N	N	7.98	nd
		A2	5-15/19	C/W	10 YR 3/2	2.5Y 5/2	SL	7.29	SBK/1/f	so/po			c/f-m	c/SFB			N	N	8.14	nd
		AC	15/19-37	G/W	10YR 4/4	10YR 6/3	S	11.67	SG/0/f	so/po			f/f	f/SFB	f/F2M		N	N	8.63	nd
		Cg	37+	U	2.5Y 4/4	2.5Y 6/3	S	3.67	SG/0/f	so/po			f/f				N	N	8.65	nd
Typic Psammaquent	TES	A1	0-5	A/W	1oYR 3/2	10YR 4/1	SL	9.06	SBK/1/m	so/po			c/f-m	f/SFB			N	N	8.42	nd
		A2	5-17/20	C/W	10YR 3/2	10YR 5/2	SL	4.09	SBK/1/m	so/po			c/f-m				N	N	8.42	nd
		AC	17/20-28	G/W	10YR 4/2	10YR 6/3	S	10.02	SG/0/f	so/po			f/f	f/SFB	f/OAF		N	N	8.61	nd
		Cg	28+	U	5Y 5/1	10YR 6/3	S	12.83	SG/0/f	so/po	c/d - 10YR 5/6		V/f			N	N	8.61	nd	
Typic Psammaquent	ITS	Oi	2.5-0	A/S	5Y 3/1	5Y 4/2	MK		nd	ss/p			m/f, c/d	m,c/RSF			Y	S/ST	7.47	7.40
		A	0-8/9	A/S	5Y 3/1	5Y 3/2	S	0.40					m/f-m							
		AC	8/9-19/20	A/W	10YR 3/1	10YR 4/2	S	2.65	SBK/1/m	so/po			m/f, c/d		c/F3M		N	N	7.98	7.35
		AC2g	19/20-30	A/W	5Y 3/2	2.5Y 6/2	S	8.44	SG/0/f	so/po			c/f	c/RFS	m2/OSF	m/FED	N	N	8.04	nd
		Cg	30+	U	2.5Y 4/2	2.5Y 6/3	S	11.32	SG/0/f	so/po			f/d				N	N	8.51	nd
Fluventic Psammowassent	SAS	A	0-5	AS	nd	nd	MK	10.83	nd	vf							Y	S/ST	nd	7.33
		C1	5-17	CW	N 2.5/0	5Y 5/2	S	11.97	nd	sf							Y	S/MD	8.62	7.49
		AC	17-60	GW	10Y 4/1	5Y 6/2		12.09	nd	nf					f/OSF	m/F3M	N	N	8.94	nd
		2C	60-84	AS	10Y 5/1	5Y 6/2	S	11.67	nd	nf				cSFB		RMX	N	N	8.85	nd
		2ACg	84-86	AW	???	???			nd	sf					m/OSF		Y	S/SL	nd	7.35
		3Cg	86-109	CW	10Y 5/1	5Y 6/2	S	8.78	nd	nf							N	N	nd	nd
		Cg	109-136+	U	N 5/0	5Y 6/1	S	8.44	nd	nf				f/OSF	m/F3M		N	N	8.60	nd
Typic Endoaquent	TES	OA	0-2	A/S	10YR 2/1	10YR 4/1	SL	2.31	GR/1/f	so/ps			m/vf	f/SFB			N	N	7.98	nd
		A1	2-4	A/S	10YR 3/1	10YR 4/1	SL	3.45	GR/1/f	so/ps			m/vf				N	N	7.88	nd
		A2	4-10	A/W	10YR4/3	10YR 5/2	LS	4.35	SBK/2/m	ss/p	f/f - 10YR 5/8		f/f				N	N	8.08	nd
		AC	10-22	C/W	10YR 4/2	2.5Y 6/3	LS	4.04	SBK/3/m	so/ps	c/d - 10YR 5/8	f/f - 2.5Y 5/1	f/f-m				N	N	8.45	nd
		Cg	22+	U	5Y 5/1	2.5 Y 6/3	LS	3.71	Massivo	s/p	c/f-10YR 5/6			m/SFB	f/OSF		N	N	8.42	nd
Typic Endoaquent	TES	Oi	1-0	A/S	2.5Y 2.5/1	2.5Y 3/2	MK	0.35	nd						m/OAF		N	N	7.55	nd
		A	0-3	A/S	7.5YR 3/3	7.5YR 4/2	MK	1.17	nd				m/vf	f/SFB		RMX	N	N	7.12	nd
		AC	3-15	C/W	2.5Y 5/3	2.5Y 6/3	LA	2.69	Massivo	s/p	c/d - 7.5YR 4/6	f/f - 2.5Y 5/1	c/f-m	f/RSB	f/OSF	c/FMM	N	N	8.13	nd
		Cg	15+	U	2.5Y 5/2	2.5Y 6/3	L	3.46	massivo	sv/pv	c/f 2.5Y 4/1						N	N	8.28	nd
Typic Endoaquent	ITS	Oi	0.5-0	A/S	10Y 3/2	2.5Y 3/2	MK	2.73	nd						m/OAF		N	N	7.84	nd
		A	0-4	A/S	7.5YR 3/3	7.5YR 4/1	MK	2.76	nd				m/vf		c/OSF		N	N	7.61	nd
		AC	4-23	C/W	5Y 5/2	5Y 6/2	LA	3.41	ABK/2/m	s/p	c/f - 5Y 4/1	f/d - 10YR 5/6	f/f	f/RSB		c/F3M	N	N	8.45	nd
		Cg	23+	U	N 3/0	5Y 6/2	LA	3.88	massivo	s/p	f/d - 5Y 5/6				c/OSF		Y	N	8.46	7.66
Typic Fluviwassent	SAS	A1	0-11	AW	10BG 2.5/1	nd	MK		nd	vf					c,m/SFB		Y	S/ST	nd	7.29
		A2	11-28	AW	10G 4/1	nd			nd	sf			c/d, m/f				Y	S/SL	7.93	7.38
		AC	28-66/67	AI	10Y 4/1	nd		3.38	nd	nf			f/f	f/RSB			N	N	8.40	nd
		1C	66/67-80	AW	10Y 4/1	nd	S	3.71	nd	nf					f/OSF	c/CLD	N	N	8.42	nd
		2C	80-110+	U	10Y 3/1	nd	S	5.37	nd	sf				f/SFB	c/OSF		N	N	8.17	nd
Typic Endoaquent	TES	OA	0-4	A/W	10YR 3/1	10YR 3/2	LS	3.03	GR/2/f	so/ps			m/f-m				N	S/SL	7.78	8.06
		A2	4-14	C/W	2.5Y 4/3	2.5Y 5/2	LS	3.20	SBK/3/m	ss/p			c/f-m	f/SFB			N	N	nd	nd
		AC	14-38	G/W	2.5Y 5/3	2.5Y 6/3	L	3.07	SBK/2/m	s/p	c/d - 10YR 5/6	f/d - 2.5Y 6/1	f/m	c/SFB			N	N	8.46	nd
		Cg	38-56+	U	5Y 5/3	2.5Y 6/3	LS	5.92	SBK/2/f-m	ss/ps	m/f - 5Y 5/6	c/d - 10YR 5/8		m/SFB			N	N	8.61	nd
Typic Endoaquent	ITS	Oi	0.5-0	AS	10Y 3/2	????	MK								m/OAF		Y	S/ST	nd	8.09
		A1	0-3	AS	10YR 3/2	10YR 4/4	MK	0.89					m/vf-f				N	N	6.96	nd
		AC	3-10	G/W	10Y 2.5/0	N 7/0	LA	3.87	massivo	s/p	f/d - 2.5Y 5/6		f/f	c/RSB; f/SFB		c/F3M	N	N	8.24	nd
		Cg	10+	U	10Y 5/1	5Y 6/2	SL	4.08	massivo	ss/ps					f/OSF		Y		8.49	7.61
Typic Endoaquent	ITS	Oi	4-0	AS	10YR 3/2	1oYR 4/1	MK	2.12					m/vf-f				N	S/SL	7.01	7.50
		A1	0-3	AS	5Y 5/1	5Y 6/2		3.71	massivo	s/p	f/d - 10YR 5/6		f/f-m		c/OSF	f/FMM	Y	N	8.29	nd
		AC	3-20	G/W	5Y 4/1	5Y 7/2		4.12	massivo	s/p						c/F3M	N	N	8.41	nd
		Cg	20+	U	N 2.5/0	5Y 6/2	S	4.89	SG/0/f	so/po					f/RSB;	f/OSF	N	N	nd	nd
Typic Fluviwassent	SAS	A1	0-3/4	AS	N 2.5/0	nd	MK			sf			m/f	m/RSB			Y	S/ST	nd	7.26
		A2	3/4-18/19	AW	2.5Y 4/2	nd		3.23		nf							Y	S/SL	8.22	7.78
		1ACg	18/19-62	AW	2.5Y 3/2	nd		1.76		nf			c/d	c/RSB		m/FMM	N	N	8.46	nd
		2ACg	62-76	AI	10Y 4/1	nd		3.60		sf			f/f	f/RSB	c2/OSF	c/FMM	N	N	8.31	nd
		Cg	76-109+	U	10Y 3/1	nd	S	4.85		nf			f/f	f/RSB	f2/OSF	f/F3M	N	N	8.39	nd

Soil System: TES = Terrestrial soils; ITS = intertidal soils; SAS = subaqueous soils --- Horizon master (Master): g = strong gleying, se = presence of sulphides --- Horizon boundary (Bound.): Distinctness: A = abrupt, C = clear, G = gradual, D = diffuse / Topography: S = smooth, W = wavy, I = irregular, U = unknown --- Mottles Quantity: f = few, c = common, m = many / Contrast: f = faint, d = distinct --- Texture Field estimation (Text.): MK = mucky, L = Loam, LS = Loamy Sand, S = sand, SL = Sandy Loam, SICL = Silty Clay Loam, SIL = Silt Loam --- Hydraulic Conductivity (HC) --- Structure (T) Type: GR = granular, ABK = angular blocky, SBK = subangular blocky, SG = single grain / (G) Grade: 0 = structureless, 1 = weak, 2 = moderate / (S) Size: vf = very fine, f = fine, m = medium --- Consistence: Stickiness: so = non-sticky, ss = slightly sticky, s = moderately sticky, sv = very sticky / Plasticity: po = non-plastic, ps = slightly plastic, p = moderately plastic, pv = very plastic / Fluidity: nf = non fluid; sl = slightly fluid; mf = moderately fluid; vf = very fluid; --- Mottles Quantity: f = few, c = common, m = many / Contrast: f = faint, d = distinct --- Roots. Quantity: vf = very few, f = few, c = common, m = many / Size: vf = very fine, f = fine, m = medium --- Biological concentrations (Bio). Quantity: f = few, c = common, m = many / (K) Kind: RSB = root sheaths, SFB = shell fragments. Other (Coats/films and redoximorphic features): Quantity: f = few, c = common, m = many / (K) Kind: CLD = clay depletion, FED = iron depletions, FEF = ferriargillans coats, F2M = reduced iron Fe²⁺ masses, FMM = iron-manganese masses, OAF = organoargillans concentrations, OSF = organic stains concentrations, RMX = reduced matrix --- H₂O₂ color change: Y = yes; N = no; Odour: (K) Kind: S = sulfurous; N = none; (I) Intensity: SL = slight; M D = moderate; ST = strong.